

Casimir Energy of AdS5 Electromagnetism and Cosmological Constant Problem

S. Ichinose

*Laboratory of Physics, School of Food and Nutritional Sciences, University of Shizuoka,
Yada 52-1, Shizuoka 422-8526, Japan*

** E-mail: ichinose@u-shizuoka-ken.ac.jp*

Casimir energy is calculated for the 5D electromagnetism in the *warped* geometry. It is compared with the *flat* case(arXiv:0801.3064). A new regularization, called *sphere lattice regularization*, is taken. It is based on the *minimal area principle* and is a *direct* realization of the geometrical approach to the *renormalization group*. The properly regularized form of Casimir energy, is expressed in a closed form. We numerically evaluate Λ (4D UV-cutoff), ω (5D bulk curvature, warp parameter) and T (extra space IR parameter) dependence of the Casimir energy. The *warp parameter* ω suffers from the *renormalization effect*. We examine the meaning of the weight function and finally reach a *new definition* of the Casimir energy where *the 4D momenta(or coordinates) are quantized* with the extra coordinate as the Euclidean time. We comment on the cosmological term at the end.

Keywords: sphere lattice; renormalization of boundary parameters.

1. Introduction

In the previous work,¹ we have examined 5D electromagnetism in the *flat* geometry. The extra space (y) is *periodic* (periodicity $2l$) and Z_2 -symmetry is taken into account. The (regularized) Casimir energy is expressed in a closed form: $E_{Cas}(\Lambda, l) = \frac{2\pi^2}{(2\pi)^4} \int_{1/l}^{\Lambda} d\tilde{p} \int_{1/\Lambda}^l dy \tilde{p}^3 W(\tilde{p}, y) F(\tilde{p}, y)$, $F(\tilde{p}, y) \equiv \int_{\tilde{p}}^{\Lambda} d\tilde{k} \frac{-3 \cosh \tilde{k}(2y-l) - 5 \cosh \tilde{k}l}{2 \sinh(\tilde{k}l)}$, where Λ is the 4D-momentum cutoff, $W(\tilde{p}, y)$ is a *weight function* to suppress the IR and UV divergences. The past approach ($W = 1$) tells us the quintic divergence (Λ^5) of the energy. This fact has been troubling us as the problem of the divergent cosmological constant in the 5D Kaluza-Klein theory.²⁻⁴

Let us review Casimir energy in 4D Electromagnetism. The electromagnetic field within two parallelly-placed (along z-direction, separation length $2l$) perfectly-conducting plates can be regarded as the sum of harmonic oscillators. For the x,y-directions, we take the periodic (periodicity $2L$) regularization. The energy of the 4D EM is given by

$$E_{4dEM} = E_{Cas} + E_{\beta}, \quad E_{Cas} = \sum \tilde{\omega}_{m_x m_y n}, \quad E_{\beta} = 2 \sum \frac{\tilde{\omega}_{m_x m_y n}}{e^{\beta \tilde{\omega}_{m_x m_y n}} - 1}, \quad (1)$$

where $\tilde{\omega}_{m_x m_y n}^2 = (\vec{\omega}_{m_x m_y n})^2 = (m_x \frac{\pi}{L})^2 + (m_y \frac{\pi}{L})^2 + (n \frac{\pi}{l})^2$.

E_β gives us Stefan-Boltzmann's law.

$$E_\beta = 2 \int_0^\infty \frac{dk 4\pi k^2}{\left(\frac{\pi}{L}\right)^2 \frac{1}{l}} \frac{\tilde{\omega}(k)}{e^{\beta \tilde{\omega}(k)} - 1} = (2L)^2 (2l) \int_0^\infty dk \mathcal{P}(\beta, k) = \frac{8}{\pi^2} \frac{L^2 l}{\beta^4} 3! \zeta(4) \quad , \quad (2)$$

where $\tilde{\omega}(k) = k$. $\mathcal{P}(\beta, k)$ is the Planck's radiation formula. The behavior of $\mathcal{P}(\beta, k)$ is graphically shown in Fig.1a. (We will see similar graphs in the following 5D case. The extra axis corresponds to the β -axis.) The peak curve of the graph is *hyperbolic*, $\beta k = \text{const.}$, in the (β, k) -plane (Wien's displacement law).




Fig. 1. (a-left) Graph of Planck's radiation formula. $\mathcal{P}(\beta, k) = \frac{1}{(ch)^3} \frac{1}{\pi^2} k^3 / (e^{\beta k} - 1)$ ($1 \leq \beta \leq 2$, $0.01 \leq k \leq 10$). (b-right) Behavior of $(-N_1/2) \tilde{p}^3 W_1(\tilde{p}, z) F^-(\tilde{p}, z)$ (elliptic suppression). $\Lambda = 20000$, $\omega = 5000$, $T = 1 \cdot 1.0001/\omega \leq z \leq 0.9999/T$, $\mu = \Lambda T/\omega \leq \tilde{p} \leq \Lambda$.

E_{Cas} is the sum of the *zero-point energy* over the all frequencies. It is *Casimir energy*. This quantity is *independent of the coupling* and is *dependent on the boundary(ies)*. E_{Cas} does not vanish for the temperature $(1/\beta)=0$. It is, however, formally *divergent*. We need a proper *regularization* for the *summation over the infinite degree of freedom* due to the continuity of the space-time. $E_{Cas}^\Lambda =$

$$\sum \tilde{\omega}_{m_x m_y n} g\left(\frac{\tilde{\omega}_{m_x m_y n}}{\Lambda}\right) = \sum \sqrt{\left(m_x \frac{\pi}{L}\right)^2 + \left(m_y \frac{\pi}{L}\right)^2 + \left(n \frac{\pi}{l}\right)^2} g\left(\frac{\tilde{\omega}_{m_x m_y n}}{\Lambda}\right), \quad (3)$$

where we introduce the *cut-off* function: $g(\omega/\Lambda) = \{ 1 \text{ for } 0 \leq \omega \leq \Lambda; 0 \text{ for } \omega > \Lambda \}$. Λ is the cut-off parameter for the absolute value of the 3D (x,y,z) momentum. We will take the limit $\Lambda \rightarrow \infty$ at an appropriate stage. Taking the reference point, $L \rightarrow \infty, L \gg l \rightarrow \infty$, from which we "measure" the energy, $E_{Cas}^{\Lambda 0} = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{dk_x dk_y}{(\pi/L)^2} \int_{-\infty}^{\infty} \frac{dk_z}{\pi/l} \sqrt{k_x^2 + k_y^2 + k_z^2} g\left(\frac{k}{\Lambda}\right)$, we finally obtain the *finite* result, $u = \frac{E_{Cas}^\Lambda - E_{Cas}^{\Lambda 0}}{(2L)^2} = \frac{\pi^2}{(2l)^3} \frac{B_4}{4!} = -\frac{\pi^2}{720} \frac{1}{(2l)^3}$, which does not depend on Λ . Especially there remains no $\log \Lambda$ divergence. This point is contrasting with the ordinary renormalization of *interacting* theories such as 4D QED and 4D YM. Hence we need *not* the renormalization of the wave-function and the parameter l . In the 5D flat geometry, the *renormalization of the boundary parameter* is necessary.¹ In the following, we examine the warped 5D case and find the another boundary parameter ω suffers from the renormalization effect.

2. Kaluza-Klein expansion approach

5D *massive* vector theory is given by

$$S_{5dV} = \int d^4x dz \sqrt{-G} \left(-\frac{1}{4} F_{MN} F^{MN} - \frac{1}{2} m^2 A^M A_M \right), F_{MN} = \partial_M A_N - \partial_N A_M,$$

$$ds^2 = \frac{1}{\omega^2 z^2} (\eta_{\mu\nu} dx^\mu dx^\nu + dz^2) = G_{MN} dX^M dX^N, G \equiv \det G_{AB} \quad (4)$$

where $M, N = 0, 1, 2, 3, 5$ (or z); $\mu, \nu = 0, 1, 2, 3$. Casimir energy is given by some integral where the (modified) Bessel functions, with the index $\nu = \sqrt{1 + \frac{m^2}{\omega^2}}$, appear. Hence the 5D EM limit is given by $\nu = 1$ ($m = 0$). We consider, however, the *imaginary* mass case $m = i\omega$ ($m^2 = -\omega^2$, $\nu = 0$). Instead of analyzing the $m^2 = -\omega^2$ of the massive vector (4), we take the 5D massive *scalar* theory on AdS_5 with $m^2 = -4\omega^2$, $\nu = \sqrt{4 + m^2/\omega^2} = 0$.

$$\mathcal{L} = \sqrt{-G} \left(-\frac{1}{2} \nabla^A \Phi \nabla_A \Phi - \frac{1}{2} m^2 \Phi^2 \right), \quad \nabla^A \nabla_A \Phi - m^2 \Phi + J = 0, \quad (5)$$

where $G \equiv \det G_{AB}$, $ds^2 = G_{AB} dX^A dX^B$. $\Phi(X) = \Phi(x^a, z)$ is the 5D scalar field. The background geometry is AdS_5 which takes the form: $(G_{AB}) = \text{diag}(\frac{1}{\omega^2 z^2} \eta_{ab}, \frac{1}{\omega^2 z^2})$, $\sqrt{-G} = \frac{1}{(\omega|z|)^5}$. The variable range of z is $-\frac{1}{T} \leq z \leq -\frac{1}{\omega}$ or $\frac{1}{\omega} \leq z \leq \frac{1}{T}$ ($-l \leq y \leq l$, $|z| = \frac{1}{\omega} e^{\omega|y|}$), where l and T are related as $\frac{1}{T} = \frac{1}{\omega} e^{\omega l}$. We take into account Z_2 symmetry: $z \leftrightarrow -z$. ω is the bulk curvature (AdS_5 parameter) and T^{-1} is the size of the extra space (infrared regularization parameter).

The Casimir energy E_{Cas} is given by $e^{-T^{-4} E_{Cas}} = \int \mathcal{D}\Phi \exp\{i \int d^5X \mathcal{L}\} =$

$$\int \mathcal{D}\Phi(X) \exp \left[i \int d^4x dz \frac{1}{(\omega|z|)^5} \frac{1}{2} \Phi \{ \omega^2 z^2 \partial_a \partial^a \Phi + (\omega|z|)^5 \frac{\partial}{\partial z} \frac{1}{(\omega z)^3} \partial_z \Phi - m^2 \Phi \} \right] \quad (6)$$

Here we introduce, instead of $\Phi(X)$, the partially (4D world only) Fourier transformed field $\Phi_p(z)$: $\Phi(X) = \int \frac{d^4p}{(2\pi)^4} e^{ipx} \Phi_p(z)$. Eq.(6) can be rewritten as

$$s(z) \equiv \frac{1}{(\omega z)^3}, \quad \hat{L}_z \equiv \frac{d}{dz} \frac{1}{(\omega z)^3} \frac{d}{dz} - \frac{m^2}{(\omega z)^5}, \quad e^{-T^{-4} E_{Cas}} = \int \mathcal{D}\Phi_p(z) \times$$

$$\exp \left[i \int \frac{d^4p}{(2\pi)^4} 2 \int_{1/\omega}^{1/T} dz \left\{ \frac{1}{2} \Phi_p(z) \left\{ -\frac{1}{(\omega z)^3} p^2 + \frac{d}{dz} \frac{1}{(\omega z)^3} \frac{d}{dz} - \frac{m^2}{(\omega z)^5} \right\} \Phi_p(z) \right\} \right] \quad (7)$$

We consider the Bessel eigen-value problem: $\{s(z)^{-1} \hat{L}_z + M_n^2\} \psi_n(z) = 0$, with Z_2 -property: $\psi_n(z) = -\psi_n(-z)$ for $P = -$; $\psi_n(z) = \psi_n(-z)$ for $P = +$, and the appropriate b.c. at fixed points. Because the set $\{\psi_n(z)\}$ constitute the orthonormal and complete system, we can express $\Phi_p(z)$ as, $\Phi_p(z) = \sum_n c_n(p) \psi_n(z)$. Eq.(7) can be further rewritten as $e^{-T^{-4} E_{Cas}} =$

$$\int \mathcal{D}\Phi_p(z) \exp \left[i \int \frac{d^4p}{(2\pi)^4} 2 \int_{1/\omega}^{1/T} dz \left\{ \frac{1}{2} \Phi_p(z) s(z) (s(z)^{-1} \hat{L}_z - p^2) \Phi_p(z) \right\} \right]$$

$$= \int \prod_n dc_n(p) \exp \left[i \int \frac{d^4p}{(2\pi)^4} \sum_n \left\{ \frac{-1}{2} c_n(p)^2 (p^2 + M_n^2) \right\} \right] = \exp \sum_{n,p} \left\{ \frac{-1}{2} \ln(p^2 + M_n^2) \right\} \quad (8)$$

where the orthonormal relation, $2 \int_{\frac{1}{\omega}}^{\frac{1}{T}} \psi_n(z) s(z) \psi_m(z) dz = \delta_{nm}$, is used. This shows that $s(z)$, defined in (7), plays the role of "inner product measure" in the function space $\{\psi_n(z), 1/\omega \leq z \leq 1/T\}$. The expression (8) is the familiar one (expanded form) of the Casimir energy.

3. Heat-Kernel Approach and Position/Momentum Propagator

Instead of the KK-expansion form, we can formally integrate out the $\Phi_p(z)$ variable in the path-integral (8).

$$\begin{aligned} e^{-T^{-4} E_{Cas}} &= \exp \left[T^{-3} \int \frac{d^4 p}{(2\pi)^4} 2 \int_{1/\omega}^{1/T} dz s(z) \left\{ -\frac{1}{2} \ln(-s(z)^{-1} \hat{L}_z + p^2) \right\} \right] \\ &= \exp \left[T^{-3} \int \frac{d^4 p}{(2\pi)^4} 2 \int_{1/\omega}^{1/T} dz s(z) \left\{ \frac{1}{2} \int_0^\infty \frac{1}{t} e^{t(s(z)^{-1} \hat{L}_z - p^2)} dt + \text{const} \right\} \right], \quad (9) \end{aligned}$$

The above formal result can be *precisely* defined using the heat equation.⁵

$$\begin{aligned} e^{-T^{-4} E_{Cas}} &= (\text{const}) \times \exp T^{-4} \int \frac{d^4 p}{(2\pi)^4} 2 \int_0^\infty \frac{1}{2} \frac{dt}{t} \text{Tr } H_p(z, z'; t), \\ \text{Tr } H_p(z, z'; t) &= \int_{1/\omega}^{1/T} s(z) H_p(z, z'; t) dz, \left\{ \frac{\partial}{\partial t} - (s^{-1} \hat{L}_z - p^2) \right\} H_p(z, z'; t) = 0. \quad (10) \end{aligned}$$

The heat kernel $H_p(z, z'; t)$ is formally solved, using the Dirac's bra and ket vectors $\langle z|, |z\rangle$, as $H_p(z, z'; t) = \langle z| e^{-(s^{-1} \hat{L}_z + p^2)t} |z'\rangle$. Using the set $\{\psi_n(z)\}$ defined previously, the explicit solution of (10) is given by

$$\begin{Bmatrix} H_p(z, z'; t) \\ E_p(z, z'; t) \end{Bmatrix} = \sum_{n \in \mathbf{Z}} e^{-(M_n^2 + p^2)t} \frac{1}{2} \{ \psi_n(z) \psi_n(z') \mp \psi_n(z) \psi_n(-z') \}, \quad P = \mp, \quad (11)$$

We here introduce the position/momentum propagators $G_p^\mp$ as follows. $G_p^\mp(z, z') \equiv$

$$\int_0^\infty dt \begin{Bmatrix} H_p(z, z'; t) \\ E_p(z, z'; t) \end{Bmatrix} = \sum_{n \in \mathbf{Z}} \frac{1}{M_n^2 + p^2} \frac{1}{2} \{ \psi_n(z) \psi_n(z') \mp \psi_n(z) \psi_n(-z') \}. \quad (12)$$

Therefore the Casimir energy E_{Cas} is given by

$$\begin{aligned} E_{Cas}(\omega, T) &= \int \frac{d^4 p}{(2\pi)^4} 2 \int_0^\infty \frac{dt}{t} 2 \int_{1/\omega}^{1/T} dz s(z) H_p(z, z; t) \\ &= \int \frac{d^4 p}{(2\pi)^4} 2 \int_0^\infty \frac{dt}{t} 2 \int_{1/\omega}^{1/T} dz s(z) \left\{ \sum_{n \in \mathbf{Z}} e^{-(M_n^2 + p^2)t} \psi_n(z)^2 \right\}, \quad (13) \end{aligned}$$

The P/M propagators $G_p^\mp$ in (12) can be expressed in a *closed* form. (See, for example, Ref. 6.) Taking the *Dirichlet* condition at all fixed points, the expression for the fundamental region $(1/\omega \leq z \leq z' \leq 1/T)$ is given by $G_p^\mp(z, z') =$

$$\mp \frac{\omega^3}{2} z^2 z'^2 \frac{\{ \mathbf{I}_0(\frac{\tilde{p}}{\omega}) \mathbf{K}_0(\tilde{p}z) \mp \mathbf{K}_0(\frac{\tilde{p}}{\omega}) \mathbf{I}_0(\tilde{p}z) \} \{ \mathbf{I}_0(\frac{\tilde{p}}{T}) \mathbf{K}_0(\tilde{p}z') \mp \mathbf{K}_0(\frac{\tilde{p}}{T}) \mathbf{I}_0(\tilde{p}z') \}}{\mathbf{I}_0(\frac{\tilde{p}}{T}) \mathbf{K}_0(\frac{\tilde{p}}{\omega}) - \mathbf{K}_0(\frac{\tilde{p}}{T}) \mathbf{I}_0(\frac{\tilde{p}}{\omega})}, \quad (14)$$

where $\tilde{p} \equiv \sqrt{p^2}$, $p^2 \geq 0$ (space-like). We can express the Λ -regularized Casimir energy in terms of the following functions $F^\mp(\tilde{p}, z)$.

$$E_{Cas}^{\Lambda, \mp}(\omega, T) = \int \frac{d^4 p}{(2\pi)^4} \Big|_{\tilde{p} \leq \Lambda} \int_{1/\omega}^{1/T} dz F^\mp(\tilde{p}, z) \quad , \quad (15)$$

where $F^\mp(\tilde{p}, z) \equiv s(z) \int_{p^2}^{\Lambda^2} \{G_k^\mp(z, z)\} dk^2 \equiv \int_{\tilde{p}}^{\Lambda} \mathcal{F}^\mp(\tilde{k}, z) d\tilde{k}$. Here we introduce the UV cut-off parameter Λ for the 4D momentum space. In Fig.2a, we show the behaviour of $\mathcal{F}^-(\tilde{k}, z)$. The table-shape graph says the "Rayley-Jeans" dominance. That is, for the wide-range region $(\tilde{p}, z)$ satisfying both $\tilde{p}(z - \frac{1}{\omega}) \gg 1$ and $\tilde{p}(\frac{1}{T} - z) \gg 1$,

$$\mathcal{F}^-(\tilde{p}, z) \approx \frac{-1}{2}, \mathcal{F}^+(\tilde{p}, z) \approx \frac{-1}{2}; \quad \tilde{p}(z - \frac{1}{\omega}) \gg 1 \text{ and } \tilde{p}(\frac{1}{T} - z) \gg 1. \quad (16)$$




Fig. 2. (a-left) Behavior of $\ln |\frac{1}{2} \mathcal{F}^-(\tilde{k}, z)| = \ln |\tilde{k} G_k^-(z, z)/(\omega z)^3|$. $\omega = 10^4, T = 1, \Lambda = 2 \times 10^4$. $1.0001/\omega \leq z \leq 0.9999/T$. $\Lambda T/\omega \leq \tilde{k} \leq \Lambda$. Note $\ln |(1/2) \times (-1/2)| \approx -1.39$. (b-right) Behaviour of $(-1/2) \tilde{p}^3 F^-(\tilde{p}, z)$ (17). $T = 1, \omega = 10^4, \Lambda = 4 \cdot 10^4$. $1.0001/\omega \leq z < 0.9999/T$, $\Lambda T/\omega \leq \tilde{p} \leq \Lambda$.

4. UV and IR Regularization Parameters and Evaluation of Casimir Energy

The integral region of the above equation (15) is displayed in Fig.3a. In the figure, we introduce the regularization cut-offs for the 4D-momentum integral, $\mu \leq \tilde{p} \leq \Lambda$. As for the extra-coordinate integral, it is the finite interval, $1/\omega \leq z \leq 1/T = e^{\omega l}/\omega$, hence we need not introduce further regularization parameters. For simplicity, we take the following IR cutoff of 4D momentum : $\mu = \Lambda \cdot \frac{T}{\omega} = \Lambda e^{-\omega l}$.

Let us evaluate the (Λ, T) -regularized value of (15).

$$E_{Cas}^{\Lambda, \mp}(\omega, T) = \frac{2\pi^2}{(2\pi)^4} \int_{\mu}^{\Lambda} d\tilde{p} \int_{1/\omega}^{1/T} dz \tilde{p}^3 F^\mp(\tilde{p}, z) \quad , \quad (17)$$

where $F^\mp(\tilde{p}, z) = \frac{2}{(\omega z)^3} \int_{\tilde{p}}^{\Lambda} \tilde{k} G_k^\mp(z, z) d\tilde{k}$. The integral region of $(\tilde{p}, z)$ is the rectangle shown in Fig.3a.




Fig. 3. (a-left) Space of $(z, \tilde{p})$ for the integration. The hyperbolic curve will be used in Sec.5. (b-right) Space of $(\tilde{p}, z)$ for the integration (present proposal).

Note that eq.(17) is the *rigorous* expression of the (Λ, T) -regularized Casimir energy. We show the behaviour of $(-1/2)\tilde{p}^3 F^-(\tilde{p}, z)$ in Fig.2b. [The requirement for the three parameters ω, T, Λ is $\Lambda \gg \omega \gg T$.] From a *close* numerical analysis of $(\tilde{p}, z)$ -integral (17), we have confirmed: $E_{Cas}^{\Lambda, -}(\omega, T) = \frac{2\pi^2}{(2\pi)^4} \times \left[-0.0250 \frac{\Lambda^5}{T} \right]$, which does *not* depend on ω . $\ln \frac{\Lambda}{T}$ -term does not appear. Compared with the flat case ($E_{Cas}^{flat}(\Lambda, l) = (1/8\pi^2) \left[-0.1249 l \Lambda^5 - (1.41, 0.706, 0.353) \times 10^{-5} l \Lambda^5 \ln(l\Lambda) \right]$), we see the factor T^{-1} plays the role of *IR cut-off* of the extra space. We note that the behavior of Fig.2b is similar to the Rayleigh-Jeans's region (small momentum region) of the Planck's radiation formula (Fig.1a) in the sense that $\tilde{p}^3 F(\tilde{p}, z) \propto \tilde{p}^3$ for $\tilde{p} \ll \Lambda$.

From the Fig.2b, the approximate form of $F(\tilde{p}, z)$ for the large Λ and $1/T$ is given by: $F^\mp(\tilde{p}, z) \approx -\frac{f}{2}\Lambda(1 - \frac{\tilde{p}}{\Lambda})$; $f = 1$, which does *not* depend on z, ω and T . f is the degree of freedom.

5. UV and IR Regularization Surfaces, Principle of Minimal Area and Renormalization Flow

In order to avoid the Λ^5 -divergence, a proposal was presented by Randall and Schwartz.⁷ They introduced the *position-dependent cut-off*, $\mu < \tilde{p} < \Lambda/\omega u$, $u \in [1/\omega, 1/T]$, for the 4D-momentum integral in the "brane" located at $z = u$. See Fig.3a. We have confirmed that the value E_{Cas} of (17), when the Randall-Schwartz integral region (Fig.3a) is taken, is proportional to Λ^5 : $E_{Cas}^{RS}(\omega, T) = \frac{2\pi^2}{(2\pi)^4} \frac{\Lambda^5}{\omega} \left\{ -1.58 \times 10^{-2} - 1.69 \times 10^{-4} \ln \frac{\Lambda}{\omega} \right\}$, which is *independent* of T . This shows the divergence situation does *not* improve compared with the non-restricted case: $E_{Cas}^{\Lambda, -}$ in Sec.4. T of $E_{Cas}^{\Lambda, -}$ is replaced by the warp parameter ω . These are contrasting with the flat case ($E_{Cas}^{RS, flat} = (1/8\pi^2)[-0.0893 \Lambda^4]$; No $\ln(l\Lambda)$ -term).

Although they claim the holography is behind the procedure, the legitimacy of the restriction looks less obvious. We have proposed an alternate approach and given a legitimate explanation within the 5D QFT.^{1,6,8,9} See Fig. 3b. The restriction region is bounded by two minimal surfaces, IR-surface and UV-surface. They are 4 dimensional manifold made from S^3 -spheres "running" along z -axis. On the 'brane' at fixed z , the 4D region bounded by IR-surface is regarded as a large-size 4D-ball and that by UV-surface is regarded as a small-size 4D-ball. Hence this regularization configuration is the 4D *sphere lattice* running along z -axis. See Ref. 10.

6. Weight Function and Casimir Energy Evaluation

We introduce, instead of restricting the integral region, a *weight function* $W(\tilde{p}, z)$ in the $(\tilde{p}, z)$ -space for the purpose of suppressing UV and IR divergences of the Casimir Energy.

$$E_{Cas}^{\mp W}(\omega, T) \equiv \int \frac{d^4 p}{(2\pi)^4} \int_{1/\omega}^{1/T} dz W(\tilde{p}, z) F^{\mp}(\tilde{p}, z), F^{\mp}(\tilde{p}, z) = \frac{2}{(\omega z)^3} \int_{\tilde{p}}^{\infty} \tilde{k} G_k^{\mp}(z, z) d\tilde{k},$$

$$\text{Examples} \begin{cases} (N_1)^{-1} e^{-(1/2)\tilde{p}^2/\omega^2 - (1/2)z^2 T^2} \equiv W_1(\tilde{p}, z), & N_1 = 1.711/8\pi^2 & \text{elliptic} \\ (N_2)^{-1} e^{-\tilde{p}zT/\omega} \equiv W_2(\tilde{p}, z), & N_2 = 2\frac{\omega^3}{T^3}/8\pi^2 & \text{hyperbolic} \\ (N_8)^{-1} e^{-1/2(\tilde{p}^2/\omega^2 + 1/z^2 T^2)} \equiv W_8(\tilde{p}, z), & N_8 = 0.4177/8\pi^2 & \text{reciprocal} \end{cases} \quad (18)$$

where $G_k^{\mp}(z, z)$ are defined in (14). N_i 's are the normalization constants. We show the shape of the energy integrand $(-1/2)\tilde{p}^3 W_1(\tilde{p}, z) F^-(\tilde{p}, z)$ in Fig.1b.

We can check the divergence (scaling) behavior of $E_{Cas}^{\mp W}$ by *numerically* evaluating the $(\tilde{p}, z)$ -integral (18) for the rectangle region of Fig.3a. $E_{Cas}^W =$

$$\begin{cases} \frac{\omega^4}{T} \Lambda \left\{ 0.42 + 0.01 \ln \frac{\Lambda}{\omega} + 0 \times \ln \frac{\Lambda}{T} \right\} & \text{for } W_1 \\ \frac{T^2}{\omega^2} \Lambda^4 (6.45 + 0.234 \ln \frac{\Lambda}{\omega} - 0.479 \ln \frac{\Lambda}{T}) \times 10^{-2} & \text{for } W_2 \\ \frac{\omega^4}{T} \Lambda \left\{ 0.42 + 0.02 \ln \frac{\Lambda}{\omega} + 0 \times \ln \frac{\Lambda}{T} \right\} & \text{for } W_8 \end{cases} \quad (19)$$

They give, after normalizing the factor Λ/T , *only* the *log-divergence*.

$$E_{Cas}^W/\Lambda T^{-1} = \alpha \omega^4 (1 - 4c \ln(\Lambda/\omega)) \quad , \quad (20)$$

where α and c can be read from (19) depending on the choice of W . We have confirmed the above result for other many examples.¹⁰ This means the 5D Casimir energy is *finitely* obtained by the ordinary renormalization of the warp factor ω . (See the final section.) In the above result of the warped case, the IR parameter l in the flat result ($E_{Cas}^{W, flat}/\Lambda l = -\frac{\alpha}{l^4} (1 - 4c \ln(l\Lambda)) = -\frac{\alpha}{l^4}$) is replaced by the inverse of the warp factor ω .

7. Meaning of Weight Function and Quantum Fluctuation of Coordinates and Momenta

In order to most naturally accomplish the above procedure, we can go to a *new step*. Namely, we *propose* to *replace* the 5D space integral with the weight W , (18), by the following *path-integral*.¹¹ We *newly define* the Casimir energy in the higher-dimensional theory as follows. $\mathcal{E}_{Cas}(\omega, T, \Lambda) \equiv$

$$\int_{\frac{1}{\Lambda}}^{\frac{1}{\mu}} d\rho \int_{r(1/\omega)=\rho}^{r(1/T)=\rho} \prod_{a,z} \mathcal{D}x^a(z) F(\frac{1}{r}, z) \exp \left[-\frac{1}{2\alpha'} \int_{1/\omega}^{1/T} \frac{1}{\omega^4 z^4} \sqrt{r'^2 + 1} r^3 dz \right], \quad (21)$$

where $\mu = \Lambda T/\omega$ and the limit $\Lambda T^{-1} \rightarrow \infty$ is taken. The string (surface) tension parameter $1/2\alpha'$ is introduced. (Note: Dimension of α' is [Length]⁴.) The square-bracket $([\dots])$ -parts of (21) are $-\frac{1}{2\alpha'} \text{Area} = -\frac{1}{2\alpha'} \int \sqrt{\det g_{ab}} d^4 x$ where g_{ab} is the induced metric on the 4D surface. $F(\tilde{p}, z)$ is defined in (15) and shows the *field-quantization* of the bulk scalar (EM) fields.

The proposed definition, (21), clearly shows the 4D space-coordinates x^a or the 4D momentum-coordinates p^a are *quantized* (quantum-statistically, not field-theoretically) with the Euclidean time z and the "area Hamiltonian" $A = \int \sqrt{\det g_{ab}} d^4 x$. [We recall

the similar situation occurs in the standard string approach. The space-time coordinates obey some uncertainty principle.¹²] Note that $F(\tilde{p}, z)$ or $F(1/r, z)$ appears, in (21), as the *energy density operator* in the quantum statistical system of $\{p^a(z)\}$ or $\{x^a(z)\}$.

In the view of the previous paragraph, the treatment of Sec.6 is an *effective* action approach using the weight function $W(\tilde{p}, z)$. Note that the integral over (p^μ, z) -space, appearing in (15), is the summation over all degrees of freedom of the 5D space(-time) points using the "naive" measure $d^4p dz$. An important point is that we have the possibility to take another measure for the summation. We have adopted, in Sec.6, the new measure $W(p^\mu, z)d^4p dz$ in such a way that the Casimir energy *does not show physical divergences*. We expect the direct evaluation of (21), numerically or analytically, leads to the similar result.

8. Discussion and Conclusion

The log-divergence, (20), is renormalized away by

$$E_{Cas}^W/\Lambda T^{-1} = \alpha \omega'^4 \quad , \quad \omega' = \omega \sqrt[4]{1 - 4c \ln(\Lambda/\omega)} \quad . \quad (22)$$

Taking into account the fact $|c| \ll 1$, we find the renormalization group function for the warp factor ω as

$$|c| \ll 1 \quad , \quad \omega' = \omega(1 - c \ln(\Lambda/\omega)) \quad , \quad \beta(\beta\text{-function}) \equiv \frac{\partial}{\partial(\ln \Lambda)} \ln \frac{\omega'}{\omega} = -c \quad . \quad (23)$$

We should notice that, in the flat geometry case, the IR parameter (extra-space size) l is renormalized. In the present warped case, however, the corresponding parameter T is *not renormalized*, but the *warp parameter* ω *is renormalized*. Depending on the sign of c , the 5D bulk curvature ω *flows* as follows. When $c > 0$, the bulk curvature ω decreases (increases) as the measurement energy scale Λ increases (decreases). When $c < 0$, the flow goes in the opposite way.

Recently the dark energy (as well as the dark matter) in the universe is a hot subject. It is well-known that the dominant candidate is the cosmological term. We also know the proto-type higher-dimensional theory, that is, the 5D KK theory, has predicted so far the *divergent* cosmological constant.⁴ This unpleasant situation has been annoying us for a long time. If we apply the present result, the situation drastically improve. The cosmological constant λ appears as

$$S = \int d^4x \sqrt{-g} \left\{ \frac{1}{G_N} (R + \lambda) \right\} + \int d^4x \sqrt{-g} \{ \mathcal{L}_{mat} \}, R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R - \lambda g_{\mu\nu} = T_{\mu\nu}^{mat}, \quad (24)$$

where G_N is the Newton's gravitational constant, R is the Riemann scalar curvature. The constant λ observationally takes the value: $\frac{1}{G_N} \lambda_{obs} \sim \frac{1}{G_N R_{cos}^2} \sim m_\nu^4 \sim (10^{-3} \text{eV})^4$, where R_{cos} is the cosmological size (Hubble length), m_ν is the neutrino mass. [The relation $m_\nu \sim \sqrt{M_{pl}/R_{cos}} = \sqrt{1/R_{cos} \sqrt{G_N}}$ appears in some extra dimension model.^{13,14} The neutrino mass is located at the geometrical average of two extreme ends of the mass scales in the universe.] On the other hand, we theoretically so far have the value: $\frac{1}{G_N} \lambda_{th} \sim \frac{1}{G_N^2} = M_{pl}^4 \sim (10^{28} \text{eV})^4$. This is because the mass scale usually comes from the quantum gravity. (See Ref. 15 for the derivation using the Coleman-Weinberg mechanism.) We have the famous huge discrepancy factor $\lambda_{th}/\lambda_{obs} \sim 10^{124}$ which appears in the Dirac's large number theory.¹⁶ If we apply the present approach, we have the warp factor ω , and the result (22) strongly suggests the following choice:

$$\frac{1}{G_N} \tilde{\lambda}_{th} = \alpha \omega'^4 \quad , \quad \omega' \sim \frac{1}{\sqrt[4]{G_N R_{cos}^2}} = \sqrt{\frac{M_{pl}}{R_{cos}}} \sim m_\nu \sim 10^{-3} \text{eV} \quad , \quad \tilde{\lambda}_{th} \sim \frac{1}{R_{cos}^2}. \quad (25)$$

Note that this choice ω' is within the allowed region obtained from the Newton force test (see Ref. 17). We succeed in obtaining the *finite* cosmological constant and its gross absolute value consistent with the observed one. Now we can understand that the *smallness of the cosmological constant comes from the renormalization flow* for the non asymptotic-free case ($c < 0$ in (23)). [In the 2 dim R^2 -gravity, the same thing occurs.¹⁸] In this case the choice of the regularization parameters are

$$T \sim 10^{-33} \text{ eV}((\text{Cosm. Size})^{-1}, 1/R_{\text{cos}}) \quad , \quad \Lambda \sim 10^{28} \text{ eV}(\text{Planck mass}, M_{pl}) \quad , \quad (26)$$

and the relations, $\omega \sim \sqrt{T\Lambda}$ and $\mu = \Lambda T/\omega \sim \sqrt{T\Lambda}$ are valid. The normalization factor in (20) is given by $\frac{\Lambda}{T} \sim 10^{61}$. The total number of the unit spheres within the large sphere is given by $\frac{\Lambda^4}{\mu^4} = \frac{\omega^4}{T^4} \sim 10^{120}$, which is expected to show the degree of freedom of the universe (4D space-time).

References

1. S. Ichinose, arXiv:0801.3064v8[hep-th], to be published in Prog.Theor.Phys.(2009).
2. Th. Kaluza, Sitzungsberichte der K.Preussischen Akademie der Wissenschaften zu Berlin. p966 (1921)
3. O. Klein, Z. Physik **37** 895 (1926)
4. T. Appelquist and A. Chodos, Phys.Rev.**D28**(1983)772
T. Appelquist and A. Chodos, Phys.Rev.Lett.**50**(1983)141
5. J. Schwinger, Phys.Rev.**82**(1951)664
6. S. Ichinose and A. Murayama, Phys.Rev.**D76**(2007)065008, hep-th/0703228
7. L. Randall and M.D. Schwartz, JHEP **0111** (2001) 003, hep-th/0108114
8. S. Ichinose, Proc. of VIII Asia-Pacific Int. Conf. on Gravitation and Astrophysics , arXiv:/0712.4043
9. S. Ichinose, Int.Jour.Mod.Phys.23A(2008)2245-2248, Proc. of Int. Conf. on Prog. of String Theory and Quantum Field Theory , arXiv:/0804.0945
10. S. Ichinose, arXiv:0812.1263[hep-th], "Casimir Energy of 5D Warped System and Sphere Lattice Regularization".
11. R.P. Feynman, "Statistical Mechanics", W.A.Benjamin,Inc., Massachusetts, 1972
12. T. Yoneya, *Duality and Indeterminacy Principle in String Theory* in "Wandering in the Fields", eds. K. Kawarabayashi and A. Ukawa (World Scientific,1987), p.419
T. Yoneya, Prog.Theor.Phys.**103**(2000)1081
13. S. Ichinose, hep-th/0012255, US-00-11, 2000
14. S. Ichinose, Proc. of 10th Tohwa Int. Symp. on String Thery, C2002, AIP Conf. Proc. 607, American Inst. Phys., Melville, New York, p307
15. S. Ichinose, Nucl.Phys.**B231**(1984)335
16. P.A.M. Dirac, Nature **139**(1937)323; Proc.Roy.Soc.**A165**(1938)199; "Directions in Physics", John Wiley & Sons, Inc., New York, 1978
17. S. Ichinose, Class.Quantum.Grav.**18**(2001)421, hep-th/0003275
18. S. Ichinose, Nucl.Phys.**B457**(1995)688